

Review

# Genetic network during neural crest induction: From cell specification to cell survival

Ben Steventon<sup>a</sup>, Carlos Carmona-Fontaine<sup>a</sup>, Roberto Mayor<sup>a,b,c,\*</sup>

<sup>a</sup> Department of Anatomy and Developmental Biology, University College London, Gower Street, London WC1E 6BT, UK

<sup>b</sup> Millennium Nucleus in Developmental Biology, Universidad de Chile, Chile

<sup>c</sup> Fundación Ciencia Para la Vida, Chile

Available online 5 August 2005

## Abstract

The concerted action of extracellular signals such as BMP, Wnt, FGF, RA and Notch activate a genetic program required to transform a naïve ectodermal cell into a neural crest cell. In this review we will analyze the extracellular signals and the network of transcription factors that are required for this transformation. We will propose the division of this complex network of factors in two main steps: an initial cell specification step followed by a maintenance or cell survival step.

© 2005 Elsevier Ltd. All rights reserved.

**Keywords:** Neural crest; Genetic cascade; Cell survival; Wnt; FGF; BMP; Retinoic acid

## Contents

1. Introduction .....	647
2. Extracellular Cell Signalling during neural crest induction .....	648
3. Early genetic network involved in Cell Specification of the neural crest .....	649
4. Specific neural crest genes involved in Cell Survival .....	650
4.1. Snail/Slug .....	650
4.2. Id3 .....	650
4.3. Sox .....	650
4.4. Twist .....	650
4.5. FoxD3 .....	651
5. Concluding remarks .....	651
Acknowledgements .....	651
References .....	651

## 1. Introduction

The neural crest is a unique and highly specialized population of cells found in all vertebrate embryos that has fascinated generations of developmental and evolutionary biologists. The neural crest develops at the border between the neural plate and the epidermis, and following closure of

the neural tube these cells delaminate from the dorsal neural tube to migrate along different pathways. On reaching their destination in the embryo, they differentiate into a wide variety of cell types.

In this review, we will analyse recent advances on the signals that induce neural crest and the genes activated by these signals that are required for the early steps of neural crest specification. Induction of neural crest is a multi-step process, and we will propose here some of the steps and the genes that control it.

\* Corresponding author. Tel.: +44 20 7679 3323; fax: +44 20 7679 7349.  
E-mail address: r.mayor@ucl.ac.uk (R. Mayor).




Fig. 1. Neural crest genetic network. Genetic network based on expression patterns and gain- and loss- of function experiments. Green square: Cell Specification genes; Blue square: Cell Survival genes. Genes written in blue indicate antiapoptotic activity. See text for more details.

Complete induction of neural crest cells requires the transformation of a naïve ectoderm into neural crest. This is a complex process that involves the activity of extracellular factors produced by the inductive tissues (*Cell Signalling* in Fig. 1), which activate a genetic program in the ectodermal cells (*Cell Specification*, green arrow in Fig. 1). Once this genetic program is activated in the neural crest cells a second set of genes, the proper neural crest genes (blue box in Fig. 1), are transcribed. One of the important functions of this second set of genes is to allow survival of the recently induced neural crest cells (maintenance or *Cell Survival* in Fig. 1). Finally, the joint action of the Cell Specification and Cell Survival genes work together to control delamination, migration and differentiation of the neural crest cells. In this review we will analyse the factors and genes involved in Cell Signalling, Specification and Survival, but we will not include the analysis of delamination, migration and differentiation as they have been discussed in another review of this issue [1].

The genetic cascade proposed in this review is an updated version of previous proposals [2,3] and it will be based in the expression pattern and gain- and loss- of function experiments performed in different animal models. Thus, although useful as a proposal is far of being definitive as not many cys-regulatory analysis has been carried out. A further limitation of this proposal is that the interaction between the genes is

not linear, and many of the genes involved in the early steps of neural crest specification are also involved in later steps of crest development.

## 2. Extracellular Cell Signalling during neural crest induction

Induction of the neural crest involves a complex set of extracellular signals that transform the fate of cells lying along the medio-lateral and anterior-posterior axes of the embryo. The signals that position the neural crest cells along these axes are released from the neural plate, the epidermis and the lateral mesoderm [4–10].

It has been shown in *Xenopus* and zebrafish embryos that prior to gastrulation BMP signalling is active throughout the entire ectoderm [11,12]. At this time, the dorsal mesoderm releases anti-BMP molecules such as noggin, follistatin and chordin, that directly bind to BMPs, inhibiting their activity. In addition, the expression of BMPs is down regulated at the most dorsal ectoderm during gastrulation by a Wnt-dependent mechanism [13]. As a consequence of all these interactions a dorso-ventral gradient of BMP activity is generated. Strong evidence in *Xenopus* and zebrafish embryos indicate that neural plate, neural crest and epidermis are specified at progressively higher levels of BMP activity [9,14–18]. The addition of BMPs to dissected neural crest induces the expression of neural crest markers in chick, supporting a role for this molecule in this animal model as well [19]. However, in this latter work it is likely that BMP is playing a role in a late step of neural crest development as the tissue used in these experiments was taken from relatively older embryos (see [20] for a comparison between the stages of different animal models concerning the timing of neural crest induction). Nevertheless, although intermediate BMP activity is required for neural crest induction in chick, *Xenopus* and zebrafish this molecule is not sufficient to induce neural plate or neural crest cells [21,22].

A second group of signals, Wnts, FGFs, and RA, are required for neural crest induction and have also been implicated in the antero-posterior patterning of the neuroectoderm [23–28]. These molecules come from the involuting endomesoderm located at the posterior part of the embryo. Their expression and inhibition are respectively required to posteriorize and anteriorize the neural tube [29–33], as well as to induce neural crest on the posterior neural folds [34–39].

Recently, the emphasis has shifted in favor of Wnt signalling as one of the key elements in neural crest induction [40]. Gain and loss of function experiments in chick, *Xenopus* and zebrafish show that Wnt signals are essential for neural crest induction. However they are not able to induce neural crest by itself in naïve ectoderm as they require working in combination with anti-BMP signals or in a previously induced neural plate [36,38,41–46]. A possible candidate to be the source of Wnt signalling in the chick embryo is the epidermis as it expresses Wnt6 at the right time to be the inducer

[43]; however, there is no experimental evidence demonstrating that *Wnt6* is required for neural crest formation. In *Xenopus* and zebrafish embryos *Wnt8* is a good candidate to be a neural crest inducer. It is expressed in the mesoderm that is able to induce neural crest, and gain and loss of function experiment show its role in neural crest induction [36,39,41,45,46].

FGF signalling is also important in neural crest induction as gain and loss of function experiments show an essential role of this molecule, and a direct, though transient, induction of some neural crest markers has been observed in naïve ectoderm treated with FGF8 [10,14,36,37]. Retinoic acid (RA) has also been implicated in neural crest induction in *Xenopus* and zebrafish embryos. Addition of RA to neuralized animal caps or expression of dominant negatives in *Xenopus* embryos, as well as analysis of zebrafish mutant indicate the requirement of this molecule for the correct specification or survival of the neural crest cells [39,47]. Moreover, it has been shown that the induction activity of these three molecules is, at least in part, related to its posteriorization activity [3,20,39]. Recently, it has been possible to dissociate the posteriorizing role of FGF and Wnt signalling from their ability to induce neural crest [48,49]. In conclusion, induction of neural crest requires a specific level of BMP and the concerted action of Wnt, FGF and RA signalling (see Fig. 1). It is likely that Wnt, FGF and RA works in the posteriorization of the neuroectoderm in an early step of neural crest specification and also as direct neural crest inducer at later steps.

Another important molecule involved in early crest specification is Notch. It has been proposed that Notch participates in neural crest specification by controlling BMP expression in chick and *Xenopus* embryos, although the exact mechanism could be different in these two species [50–54]. In addition, Notch signalling seems to control neural crest development by repressing neurogenesis [51]. We will not discuss Notch signalling further as it is analyzed elsewhere in this issue [55].

### 3. Early genetic network involved in Cell Specification of the neural crest

As consequence of the extracellular signalling working on the ectoderm a first set of genes that encode for transcription factors are activated in the prospective neural crest cells. These are the genes that control Cell Specification (see Fig. 1).

Gene expression pattern suggests that *Msx*, *AP2α* and *Dlx* genes are amongst the earliest genes activated in the ectoderm fated to become neural crest. We have grouped these three genes together because they exhibit a similar pattern of expression. They are initially expressed in the non-neural ectoderm, and later they are restricted to the neural fold region in *Xenopus* and chick embryos [56–58]. Interestingly amphioxus, a non vertebrate chordate, shows a similar early AP2 expression pattern suggesting that this molecule

also had an important role in neural crest evolution as well as in other vertebrate traits [59,60]. Their expression pattern correlates with the assumed ventro-dorso gradient of BMP activity, and identification of cis-regulatory elements in some of these genes indicate that is very likely to be direct targets of BMP [58,61–63]. Loss of function experiments in mouse, zebrafish and *Xenopus* embryos show that they are required for the early specification of the neural crest [44,58,59,64–67]. *Msx1* seems to play a key role in neural crest development, and epistatic experiments in *Xenopus* suggest that works upstream of the *Snail* genes [44,48,67]. However, once the neural crest is induced *Msx1* expression is switched off from the neural crest, and become to play a role in controlling neural crest apoptosis [48,68]. The zebrafish *AP2a* mutant, *mont blanc*, shows a defect in the early specification of the neural crest, as well as in neural crest derivatives [64,69–71]. It also shows a later role as an inhibitor of apoptosis in the neural crest cells [69]. Morpholino mediated knockdown experiments in *Xenopus* indicates that *AP2a* is essential for the early specification of neural crest cells [66]. Thus, *AP2a* seems to be a gene involved in early as well as late neural crest development.

*Dlx* genes are important in the positioning of the neural plate border, including neural crest and placodal cells [72–74], although *Dlx5* seems to be more specific for the neural crest cells [75]. It is important to note that *Msx*, *Dlx* and *AP2α* have an earlier role in epidermal development, and in consequence it could be suggested that specification of the neural crest at this early stage involves the dorsalization of epidermis.

The next group of genes, *Zic*, *Pax3* and *c-Myc*, are expressed in a more restricted domain that the previous one, they are not expressed in the entire epidermis at earlier stages but they are present in a wider domain than the premigratory neural crest cells [76–81]. Multiple *Zic* genes have been identified in mouse and *Xenopus* that are expressed in overlapping but distinct patterns [80,82]. Disruption of mouse *Zic* genes leads to multiple defects in neural and neural crest derivatives [82–84]. In *Xenopus*, expression of the *Zic* genes are a very early response to neural inducing signals, and ectopic expression of these *Zic* genes leads to neural plate and neural crest formation [67,78–80,85–87].

*Pax3* is expressed in the neural folds and loss of function experiments in mouse and *Xenopus* show a key role in neural crest development [48,67,88–90]. Using a combination of overexpression and morpholino mediated knockdown experiments, Monsoro-Burq et al. [48], show that *Msx1* and *Pax3* are required for neural crest formation and that *Msx1* seems to work upstream of *Pax3* (see Fig. 1). Interestingly, it is proposed in their work that Wnt controls neural crest induction through *Pax3* activity, while FGF8 requires *Msx1* and *Pax3* actions; however the molecular mechanism by which the combination of the anti-BMP molecule noggin and *Msx1* is able to induce neural crest in naïve ectoderm is not explained [48]. A similar strategy based on morpholino injections has been recently used to show a cooperative function of *Pax3*

and *Zic1* in neural crest specification and a mutual regulation between these two factors has also been shown (see Fig. 1 and [67]). The exact pathway network connecting *Msx1*, *Pax3* and the *Zic* genes requires further careful consideration.

The proto-oncogene c-Myc has been found to be expressed in the neural plate border of amphibian embryos earlier and in a broader domain than many specific neural crest genes [81]. Morpholino knockdown and overexpression experiments show an essential role of c-Myc in neural crest development that seems to be independent upon cell proliferation or cell death [81]. c-Myc is expressed almost at the same time and in the same cells as *Msx1* and *Pax3*, and it is likely that an interaction between these factors is required for the early step of neural crest specification.

#### 4. Specific neural crest genes involved in Cell Survival

Following the initial specification of neural crest precursors at the neural plate border, a network of genes is set up that function to maintain these precursors (blue arrow in Fig. 1). The genes transcribed at this step of neural crest induction are expressed later than the Cell Specification genes and only in the neural crest territory. They encode for transcription factors of the *Snail* and *Sox* family of genes, and for the genes *FoxD3*, *Id3* and *Twist*. Functional experiments are insufficient to predict a cascade of genetic interactions amongst these genes as the up or down regulation of one gene will effect the expression of others. It is also important to note that *Snail*, *Slug* and *FoxD3* all function as transcriptional repressors [91–94]; this means that there are additional transcription factors mediating the activity of these genes. In addition, as many of these genes are involved in cell survival it is not known in many cases whether the observed effect is due to changes in rates of cell death/proliferation or changes in the transcription of neural crest genes. Therefore the grey arrows included in Fig. 1 that connect these genes are tentative.

##### 4.1. *Snail/Slug*

Two members of the *Snail* family of genes, *Snail* and *Slug*, play key roles in neural crest development. The observed function of these genes appears to be dependent upon the species examined. In *Xenopus* *snail* is expressed in the prospective neural folds slightly earlier than *Slug*, and based on animal caps and epistatic experiments *Snail* seems to work upstream of *Slug* in neural crest development (see Fig. 1 and [94,95]). In addition, *Slug* can produce an expansion in the *Slug* and *Twist* expression domains in whole embryos [36]. Loss of function experiments in *Xenopus* using different dominant negative constructs have shown that *Slug* is required for full expression of *Slug*, *Sox9*, *Twist*, *Ets-1*, *FoxD3* and *Sox10* [91,92,94,96,97]. Electroporation of *Slug* into the cephalic regions of chick neural tubes produces an expansion in the

expression of several neural crest markers including HNK-1, RhoB and *Pax3* [98] but in the trunk additional signals are required [99]. In conclusion, *Snail/Slug* genes are required for the specification of the neural crest cells in chick, mouse and *Xenopus* embryos.

In addition to this early role of the *Snail* genes another important function is to control cell-cycle progression in neural crest cells [100] and to inhibit apoptosis in the crest cells, by controlling the expression of Bcl-x<sub>L</sub> and specific caspases [68,100]. Thus, this is the first example of this group of genes expressed specifically in the neural crest that have an anti-apoptotic or cell survival activity.

##### 4.2. *Id3*

Two recent studies in *Xenopus* have analysed the role of *Id3* in neural crest formation [101,102]. They showed that depletion of *Id3* by morpholino knockdown experiments results in the down-regulation of early neural crest markers (*Slug*, *Sox10*, *FoxD3* and *Twist*) and an increase in apoptosis in the neural crest cells. The down-regulation of neural crest markers in *Id3* depleted embryos occurs very early and could not be rescued by *Slug* expression suggesting that *Id3* functions to maintain the expression of neural crest progenitor genes as well as control cell proliferation and apoptosis [101,102].

##### 4.3. *Sox*

The *Sox* genes are a group of transcription factors with considerable importance in neural crest development; their role has been considered in detail elsewhere in this issue [103] and so will not be dealt with in detail here. Briefly, *Sox8*, *Sox9*, *Sox10* and *LSox5* are expressed in the neural crest of mouse, chick, zebrafish and *Xenopus* embryos, at different times during development. The most studied *Sox* genes in the neural crest are *Sox9* and *Sox10*, and multiple interactions of these genes with other members of the Cell Survival group of genes have been described (see Fig. 1 and [96,97,99,104–112]). *Sox9*-null mice mutants showed massive cell death in the trunk neural crest population prior to or shortly after delamination [99] and in *Sox10* mutant embryos the neural crest undergo apoptosis before they can differentiate [113–115]. Similarly, neural crest of the *Sox10* zebrafish mutant (*colourless*) fails to migrate and undergo cell death [107]. Taken together these results point towards a role of *Sox9* and *Sox10* as a survival factor.

##### 4.4. *Twist*

*Twist* is another gene expressed in pre-migratory neural crest cells in *Xenopus* [95] and mice [116,117]. Mice lacking functional *Twist* have defects in neural tube closure and neural crest cell migration [117]. Interestingly, *Twist* has been implicated in controlling cell proliferation and survival during mouse paraxial mesoderm development [118]. The role



of *Twist* in pre-migratory neural crest cells has not yet been examined.

#### 4.5. *FoxD3*

*FoxD3* is one of the earliest neural crest genes to be expressed in mice [119], zebrafish [120], *Xenopus* [92,93] and chicks [121,122]. Functional experiments in *Xenopus* have been somewhat contradictory. Overexpression has been reported to decrease the expression of neural crest markers while causing an expansion of the neural plate [95]. Sasai et al. [92] found that injection of *FoxD3* RNA led to the ectopic induction of *Slug*, *AP-2*, *FoxD3*, *Ets-1*, *Twist* and *Sox2* in embryos and animal caps. It is not known whether *FoxD3* can induce neural crest cells directly or whether the induction is a secondary effect of neural induction. The differences between these two studies are likely to be due to the different doses of RNA used. Indeed, injecting low doses of *FoxD3* RNA results in the expansion of neural crest markers whereas high doses result in the inhibition of neural crest markers (Francisco Romero and R.M. unpublished). Electroporation of *FoxD3* into the chick neural tube can induce HNK-1 expression in neuroepithelial cells as well as an increase in migratory neural crest cells from the dorsal neural tube [99,121,122]. *FoxD3* can also induce changes in the expression of cell adhesion molecules required for delamination, although a dominant negative of SoxE genes can inhibit the effects on HNK-1 induction, it has no effect on the cell adhesion changes [99]. It has not yet been assessed whether *FoxD3* is involved in the survival of neural crest cells.

#### 5. Concluding remarks

The combination of an intermediate level of BMP signalling together with the action of Wnt, FGF, RA and Notch activates a genetic program in the ectoderm that defines the prospective neural crest cells. Some of the genes of this early neural crest network are *Msx*, *Dlx*, *Ap2*, *Pax3*, *Zic* and *cMyc* (green box in Fig. 1). The individual and/or concerted action of these Cell Specification genes activates a second group of genes. In this second group the genes are expressed specifically in the neural crest cells and multiple interactions among each gene have been described (blue box in Fig. 1). One of the most interesting outcomes of this analysis is that for almost all the genes in this group an anti-apoptotic or cell survival activity has been described. This is why we propose to call these genes Cell Survival genes and to call this step of neural crest development the Maintenance step (Fig. 1). The only two exceptions to this observation are the *Ap2 $\alpha$*  gene (included in the Cell Specification step) which also has an anti-apoptotic activity and the *FoxD3* gene, where no analysis of its role on cell survival has been performed. Interestingly, a recent review proposes the *Ap2 $\alpha$*  gene in the same group as *Snail*, *Slug* and *Sox* genes [123].

The need for a genetic program of cell survival is not surprising, as the neural crest are likely to be induced in an hostile environment surrounded by epidermal and neural plate cells, and later they will migrate very far from its original niche. Thus, many of the Cell Specification and Cell Survival genes used during the induction of the neural crest cell will also be used during the last step of the neural crest journey: migration and differentiation.

#### Acknowledgements

We thank Lorena Marchant for comment on the manuscript. This investigation was supported by an International Research Scholar Award from the Howard Hughes Medical Institute to R.M., and by grants from MRC and the Millennium Program (P99-137F and ICM P02-050). B.S. has a BBSRC Ph.D. fellowship.

#### References

- [1] Morales AV, Barbas JA, Nieto MA. How to become a neural crest: from segregation to delamination. *Sem Cell Dev Biol* 2005;16:655–62.
- [2] Mayor R, Young R, Vargas A. Development of neural crest in *Xenopus*. *Curr Top Dev Biol* 1999;43:85–113.
- [3] Mayor R, Aybar M. Induction and development of neural crest in *Xenopus laevis*. *Cell Tissue Res* 2001;305:203–9.
- [4] Raven CP, Kloos J. Induction by medial and lateral pieces of the archenteron roof, with special reference to the determination of the neural crest. *Acta Neerl Morphol Morm Pathol* 1945;5:348–62.
- [5] Moury JD, Jacobson AG. Neural fold formation at newly created boundaries between neural plate and epidermis in the Axolotl. *Dev Biol* 1989;133:44–57.
- [6] Selleck MAJ, Bronnerfraser M. Origins of the avian neural crest—the role of neural plate–epidermal interactions. *Development* 1995;121:525–38.
- [7] Mancilla A, Mayor R. Neural crest formation in *Xenopus laevis*: mechanisms of *Xslug* induction. *Dev Biol* 1996;177:580–9.
- [8] Bonstein L, Elias S, Frank D. Paraxial-fated mesoderm is required for neural crest induction in *Xenopus* embryos. *Dev Biol* 1998;193:156.
- [9] Marchant L, Linker C, Ruiz P, Guerrero N, Mayor R. The inductive properties of mesoderm suggest that the neural crest cells are specified by a BMP gradient. *Dev Biol* 1998;198:319–29.
- [10] Monsoro-Burq AH, Fletcher RB, Harland RM. Neural crest induction by paraxial mesoderm in *Xenopus* embryos requires FGF signals. *Development* 2003;130:3111–24.
- [11] Fainsod A, Steinbeisser H, Derobertis EM. On the function of *Bmp-4* in patterning the marginal zone of the *Xenopus* embryo. *EMBO J* 1994;13:5015–25.
- [12] Hemmati-Brivanlou A, Thomsen GH. Ventral mesodermal patterning in *Xenopus* embryos—expression patterns and activities of *Bmp-2* and *Bmp-4*. *Dev Genet* 1995;17:78–89.
- [13] Baker JC, Beddington RSP, Harland RM. Wnt signaling in *Xenopus* embryos inhibits *Bmp4* expression and activates neural development. *Genes Dev* 1999;13:3149–59.
- [14] Mayor R, Morgan R, Sargent MG. Induction of the prospective neural crest of *Xenopus*. *Development* 1995;121:767–77.
- [15] Wilson PA, Lagna G, Suzuki A, Hemmati-Brivanlou A. Concentration-dependent patterning of the *Xenopus* ectoderm

- by BMP4 and its signal transducer smad1. *Development* 1997;124:3177–84.
- [16] Neave B, Holder N, Patient R. A graded response to BMP-4 spatially coordinates patterning of the mesoderm and ectoderm in the zebrafish. *Mech Dev* 1997;62:183–95.
- [17] Nguyen VH, Schmid B, Trout J, Connors SA, Ekker M, Mullins MC. Ventral and lateral regions of the zebrafish gastrula, including the neural crest progenitors, are established by a *bmp2b*/swirl pathway of genes. *Dev Biol* 1998;199:93–110.
- [18] Glavic A, Silva F, Aybar MJ, Bastidas F, Mayor R. Interplay between Notch signaling and the homeoprotein *Xiro1* is required for neural crest induction in *Xenopus* embryos. *Development* 2004;131:347–59.
- [19] Liem KF, Tremml G, Roelink H, Jessell TM. Dorsal differentiation of neural plate cells induced by Bmp-mediated signals from epidermal ectoderm. *Cell* 1995;82:969–79.
- [20] Aybar MJ, Mayor R. Early induction of neural crest cells: lessons learned from frog, fish and chick. *Curr Opin Genet Dev* 2002;12:452–8.
- [21] Linker C, Stern CD. Neural induction requires BMP inhibition only as a late step, and involves signals other than FGF and Wnt antagonists. *Development* 2004;131:5671–81.
- [22] Delaune E, Lemaire P, Kodjabachian L. Neural induction in *Xenopus* requires early FGF signalling in addition to BMP inhibition. *Development* 2005;132:299–310.
- [23] Darken RS, Wilson PA. Axis induction by Wnt Signaling: target promoter responsiveness regulates competence. *Dev Biol* 2001;234:42.
- [24] Fredieu JR, Cui Y, Maier D, Danilchik MV, Christian JL. *Xwnt-8* and lithium can act upon either dorsal mesodermal or neuroectodermal cells to cause a loss of forebrain in *Xenopus* embryos. *Dev Biol* 1997;186:100.
- [25] Glinka A, Wu W, Onichtchouk D, Blumenstock C, Niehrs C. Head induction by simultaneous repression of *Bmp* and *Wnt* signalling in *Xenopus*. *Nature* 1997;389:517.
- [26] Kiecker C, Niehrs C. A morphogen gradient of *Wnt/β*-catenin signalling regulates anteroposterior neural patterning in *Xenopus*. *Development* 2001;128:4189–201.
- [27] Kudoh T, Wilson SW, Dawid IB. Distinct roles for *Fgf*, *Wnt* and retinoic acid in posteriorizing the neural ectoderm. *Development* 2002;129:4335–46.
- [28] McGrew LL, Lai C-J, Moon RT. Specification of the anteroposterior neural axis through synergistic interaction of the Wnt Signaling Cascade with *noggin* and *follistatin*. *Dev Biol* 1995;172:337.
- [29] Blumberg B, Bolado J, Moreno TA, Kintner C, Evans RM, Papalopulu N. An essential role for retinoid signaling in anteroposterior neural patterning. *Development* 1997;124:373–9.
- [30] Cox WG, Hemmati-Brivanlou A. Caudalization of neural fate by tissue recombination and *bFGF*. *Development* 1995;121:4349–58.
- [31] Houart C, Westerfield M, Wilson SW. A small population of anterior cells patterns for forebrain during zebrafish gastrulation. *Nature* 1998;391:788–92.
- [32] Houart C, Caneparo L, Heisenberg CP, Barth KA, Take-Uchi M, Wilson SW. Establishment of the telencephalon during gastrulation by local antagonism of *Wnt* signaling. *Neuron* 2002;35:255–65.
- [33] Kazanskaya O, Glinka A, Niehrs C. The role of *Xenopus dickkopf1* in prechordal plate specification and neural patterning. *Development* 2000;127:4981–92.
- [34] Chang C, Hemmati-Brivanlou A. Neural crest induction by *Xwnt7B* in *Xenopus*. *Dev Biol* 1998;194:129.
- [35] Kengaku M, Okamoto H. Basic fibroblast growth factor induces differentiation of neural tube and neural crest lineages of cultured ectoderm cells from *Xenopus* gastrula. *Development* 1993;119:1067–78.
- [36] LaBonne C, Bronner-Fraser M. Neural crest induction in *Xenopus*: evidence for a two-signal model. *Development* 1998;125:2403–14.
- [37] Mayor R, Guerrero N, Martinez C. Role of FGF and *noggin* in neural crest induction. *Dev Biol* 1997;189:1–12.
- [38] Saint-Jeannet JP, He X, Varmus HE, Dawid IB. Regulation of dorsal fate in the neuraxis by *Wnt-1* and *Wnt-3a*. *Proc Natl Acad Sci USA* 1997;94:13713–8.
- [39] Villanueva S, Glavic A, Ruiz P, Mayor R. Posteriorization by FGF, *Wnt*, and retinoic acid is required for neural crest induction. *Dev Biol* 2002;241:289–301.
- [40] Trainor P, Krumlauf R. Development: riding the crest of the *Wnt* signaling wave. *Science* 2002;297:781–3.
- [41] Lekven AC, Thorpe CJ, Waxman JS, Moon RT. Zebrafish *wnt8* encodes two *wnt8* proteins on a bicistronic transcript and is required for mesoderm and neuroectoderm patterning. *Dev Cell* 2001;1:103–14.
- [42] Dearnoff MA, Tan C, Saint-Jeannet JP, Klein PS. A role for *frizzled 3* in neural crest development. *Development* 2001;128:3655–63.
- [43] Garcia-Castro MI, Marcelle C, Bronner-Fraser M. Ectodermal *Wnt* function as a neural crest inducer. *Science* 2002;297:848–51.
- [44] Tribulo C, Aybar MJ, Nguyen VH, Mullins MC, Mayor R. Regulation of *Msx* genes by a *Bmp* gradient is essential for neural crest specification. *Development* 2003;130:6441–52.
- [45] Lewis JL, Bonner J, Modrell M, Ragland JW, Moon RT, Dorsky RI, et al. Reiterated *Wnt* signaling during zebrafish neural crest development. *Development* 2004;131:1299–308.
- [46] Bastidas F, De Calisto J, Mayor R. Identification of neural crest competence territory: Role of *Wnt* signaling. *Dev Dyn* 2004;229:109–17.
- [47] Begemann G, Schilling TF, Rauch GJ, Geisler R, Ingham PW. The zebrafish neckless mutation reveals a requirement for *raldh2* in mesodermal signals that pattern the hindbrain. *Development* 2001;128:3081–94.
- [48] Monsoro-Burq AH, Wang E, Harland R. *Msx1* and *Pax3* cooperate to mediate FGF8 and *WNT* signals during *Xenopus* neural crest induction. *Dev Cell* 2005;8:167–78.
- [49] Wu JL, Yang J, Klein PS. Neural crest induction by the canonical *Wnt* pathway can be dissociated from anterior-posterior neural patterning in *Xenopus*. *Dev Biol* 2005;279:220–32.
- [50] Cornell RA, Eisen JS. *Delta* signaling mediates segregation of neural crest and spinal sensory neurons from zebrafish lateral neural plate. *Development* 2000;127:2873–82.
- [51] Cornell RA, Eisen JS. *Delta/Notch* signaling promotes formation of zebrafish neural crest by repressing *Neurogenin 1* function. *Development* 2002;129:2639–48.
- [52] Endo Y, Osumi N, Wakamatsu Y. Bimodal functions of *Notch*-mediated signaling are involved in neural crest formation during avian ectoderm development. *Development* 2002;129:863–73.
- [53] Endo Y, Osumi N, Wakamatsu Y. *Deltex/Dtx* mediates *NOTCH* signaling in regulation of *Bmp4* expression in cranial neural crest formation during avian development. *Dev Growth Differ* 2003;45:241–8.
- [54] Glavic A, Honore SM, Feijoo CG, Bastidas F, Allende ML, Mayor R. Role of BMP signaling and the homeoprotein *iroquois* in the specification of the cranial placodal field. *Dev Biol* 2004;272:89–103.
- [55] Cornell RA, Eisen JS. *Notch* in the pathway: the roles of *Notch* signalling in neural crest development. *Sem Cell Dev Biol* 2005;16:663–72.
- [56] Luo T, Matsuo-Takasaki M, Thomas ML, Weeks DL, Sargent TD. Transcription factor *AP-2* is an essential and direct regulator of epidermal development in *Xenopus*. *Dev Biol* 2002;245:136–44.
- [57] Davidson D. The function and evolution of *Msx* genes—pointers and paradoxes. *Trends Genet* 1995;11:405–11.
- [58] Suzuki A, Ueno N, Hemmati-Brivanlou A. *Xenopus msx1* mediates epidermal induction and neural inhibition by *BMP4*. *Development* 1997;124:3037–44.

- [59] Meulemans D, Bronner-Fraser M. Amphioxus and lamprey AP-2 genes: implications for neural crest evolution and migration patterns. *Development* 2002;129:4953–62.
- [60] Meulemans D, Bronner-Fraser M. Central role of gene cooption in neural crest evolution. *J Exp Zool B Mol Dev Evol*; 2005, May 3 [Epub ahead of print].
- [61] Park GT, Morasso MI. Bone morphogenetic protein-2 (BMP-2) transactivates Dlx3 through Smad1 and Smad4: alternative mode for Dlx3 induction in mouse keratinocytes. *Nucleic Acids Res* 2002;30:515–22.
- [62] Alvarez-Martinez CE, Binato R, Gonzalez S, Pereira M, Robert B, Abdelhay E. Characterization of a Smad Motif Similar to Drosophila Mad in the Mouse Msx 1 Promoter. *Biochem Biophys Res Commun* 2002;291:655.
- [63] Holzschuh J, Barrallo-Gimeno A, Ettl AK, Durr K, Knapik EW, Driever W. Noradrenergic neurons in the zebrafish hind-brain are induced by retinoic acid and require tfap2a for expression of the neurotransmitter phenotype. *Development* 2003;130:5741–54.
- [64] Knight RD, Nair S, Nelson SS, Afshar A, Javidan Y, Geisler R, et al. lockjaw encodes a zebrafish tfap2a required for early neural crest development. *Development* 2003;130:5755–68.
- [65] Brewer S, Feng WG, Huang J, Sullivan S, Williams T. Wnt1-Cre-mediated deletion of AP-2 alpha causes multiple neural crest-related defects. *Dev Biol* 2004;267:135–52.
- [66] Luo T, Lee YH, Saint-Jeannet JP, Sargent TD. Induction of neural crest in Xenopus by transcription factor AP2 alpha. *Proc Natl Acad Sci USA* 2003;100:527–32.
- [67] Sato TSN, Sasai Y. Neural crest determination by co-activation of Pax3 and Zic1 genes in Xenopus ectoderm. *Development* 2005;132:2355–63.
- [68] Tribulo C, Aybar MJ, Sanchez SS, Mayor R. A balance between the anti-apoptotic activity of Slug and the apoptotic activity of msx1 is required for the proper development of the neural crest. *Dev Biol* 2004;275:325–42.
- [69] Knight RD, Javidan Y, Nelson S, Zhang TL, Schilling T. Skeletal and pigment cell defects in the lockjaw mutant reveal multiple roles for zebrafish tfap2a in neural crest development. *Dev Dyn* 2004;229:87–98.
- [70] Barrallo-Gimeno A, Holzschuh J, Driever W, Knapik EW. Neural crest survival and differentiation in zebrafish depends on mont blanc/tfap2a gene function. *Development* 2004;131:1463–77.
- [71] O'Brien EK, d'Alencon C, Bonde G, Li W, Schoenebeck J, Allende ML, et al. Transcription factor Ap-2 alpha is necessary for development of embryonic melanophores, autonomic neurons and pharyngeal skeleton in zebrafish. *Dev Biol* 2004;265:246–61.
- [72] McLarren KW, Litsiou A, Streit A. DLX5 positions the neural crest and preplacode region at the border of the neural plate. *Dev Biol* 2003;259:34–47.
- [73] Woda JM, Pastagia J, Mercola M, Artinger KB. Dix proteins position the neural plate border and determine adjacent cell fates. *Development* 2003;130:331–42.
- [74] Kaji T, Artinger KB. dlx3b and dlx4b function in the development of Rohon-Beard sensory neurons and trigeminal placode in the zebrafish neurula. *Dev Biol* 2004;276:523–40.
- [75] Luo T, Matsuo-Takasaki M, Sargent TD. Distinct roles for distal-less genes Dlx3 and Dlx5 in regulating ectodermal development in Xenopus. *Mol Reprod Dev* 2001;60:331–7.
- [76] Bang AG, Papalopulu N, Kintner C, Goulding MD. Expression of Pax-3 is initiated in the early neural plate by posteriorizing signals produced by the organizer and by posterior non-axial mesoderm. *Development* 1997;124:2075–85.
- [77] Bang AG, Papalopulu N, Goulding MD, Kintner C. Expression of Pax-3 in the lateral neural plate is dependent on a Wnt-mediated signal from posterior nonaxial mesoderm. *Dev Biol* 1999;212:366–80.
- [78] Nakata K, Nagai T, Aruga J, Mikoshiba K. Xenopus Zic3 a primary regulator both in neural and neural crest development. *Proc Natl Acad Sci USA* 1997;94:11980–5.
- [79] Nakata K, Nagai T, Aruga J, Mikoshiba K. Xenopus Zic family and its role in neural and neural crest development. *Mech Dev* 1998;75:43–51.
- [80] Nakata K, Koyabu Y, Aruga J, Mikoshiba K. A novel member of the Xenopus Zic family, Zic5, mediates neural crest development. *Mech Dev* 2000;99:83–91.
- [81] Bellmeyer A, Kruse J, Lindgren J, LaBonne C. The protooncogene c-Myc is an essential regulator of neural crest formation in Xenopus. *Dev Cell* 2003;4:827–39.
- [82] Nagai T, Aruga J, Minowa O, Sugimoto T, Ohno Y, Noda T, et al. Zic2 regulates the kinetics of neurulation. *Proc Natl Acad Sci USA* 2000;97:1618–23.
- [83] Aruga J, Minowa O, Yaginuma H, Kuno J, Nagai T, Noda T, et al. Mouse Zic1 is involved in cerebellar development. *J Neurosci* 1998;18:284–93.
- [84] Inoue T, Hatayama M, Tohmonda T, Itohara S, Aruga J, Mikoshiba K. Mouse Zic5 deficiency results in neural tube defects and hypoplasia of cephalic neural crest derivatives. *Dev Biol* 2004;270:146–62.
- [85] Brewster R, Lee J, Altaba ARI. Gli/Zic factors pattern the neural plate by defining domains of cell differentiation. *Nature* 1998;393:579–83.
- [86] Kuo JS, Patel M, Gamse J, Merzdorf C, Liu XD, Apekin V, et al. opl: a zinc finger protein that regulates neural determination and patterning in Xenopus. *Development* 1998;125:2867–82.
- [87] Mizuseki K, Kishi M, Matsui M, Nakanishi S, Sasai Y. Xenopus Zic-related-1 and Sox-2, two factors induced by chordin, have distinct activities in the initiation of neural induction. *Development* 1998;125:579–87.
- [88] Borycki AG, Li J, Jin FZ, Emerson CP, Epstein JA. Pax3 functions in cell survival and in pax7 regulation. *Development* 1999;126:1665–74.
- [89] Mansouri A, Pla P, Larue L, Gruss P. Pax3 acts cell autonomously in the neural tube and somites by controlling cell surface properties. *Development* 2001;128:1995–2005.
- [90] Wiggan ON, Fadel MP, Hamel PA. Pax3 induces cell aggregation and regulates phenotypic mesenchymal-epithelial interconversion. *J Cell Sci* 2002;115:517–29.
- [91] LaBonne C, Bronner-Fraser M. Snail-related transcriptional repressors are required in Xenopus for both the induction of the neural crest and its subsequent migration. *Dev Biol* 2000;221:195–205.
- [92] Sasai N, Mizuseki K, Sasai Y. Requirement of FoxD3-class signaling for neural crest determination in Xenopus. *Development* 2001;128:2525–36.
- [93] Pohl BS, Knochel W. Overexpression of the transcriptional repressor FoxD3 prevents neural crest formation in Xenopus embryos. *Mech Dev* 2001;103:93.
- [94] Aybar MJ, Nieto MA, Mayor R. Snail precedes Slug in the genetic cascade required for the specification and migration of the Xenopus neural crest. *Development* 2003;130:483–94.
- [95] Linker C, Bronner-Fraser M, Mayor R. Relationship between gene expression domains of Xsnail, Xslug, and Xtwist and cell movement in the prospective neural crest of Xenopus. *Dev Biol* 2000;224:215–25.
- [96] Honore SM, Aybar MJ, Mayor R. Sox10 is required for the early development of the prospective neural crest in Xenopus embryos. *Dev Biol* 2003;260:79–96.
- [97] Lee YH, Aoki Y, Hong CS, Saint-Germain N, Credidio C, Saint-Jeannet JP. Early requirement of the transcriptional activator Sox9 for neural crest specification in Xenopus. *Dev Biol* 2004;275:93–103.
- [98] del Barrio MG, Nieto MA. Overexpression of Snail family members highlights their ability to promote chick neural crest formation. *Development* 2002;129:1583–93.

- [99] Cheung M, Chaboissier MC, Mynett A, Hirst E, Schedl A, Briscoe J. The transcriptional control of trunk neural crest induction, survival, and delamination. *Dev Cell* 2005;8:179–92.
- [100] Vega S, Morales AV, Ocana OH, Valdes F, Fabregat I, Nieto MA. Snail blocks the cell cycle and confers resistance to cell death. *Genes Dev* 2004;18:1131–43.
- [101] Light W, Vernon A, Lasorella A, Iavarone A, Labonne C. *Xenopus* Id3 is required downstream of Myc for the formation of multipotent neural crest progenitor cells. *Development* 2005;132:1831–41.
- [102] Kee Y, Bronner-Fraser M. To proliferate or to die: role of Id3 in cell cycle progression and survival of neural crest progenitors. *Genes Dev* 2005;19:744–55.
- [103] Hong CS, Saint-Jeannet JP. Sox proteins and neural crest development. *Sem Cell Dev Biol* 2005;16:694–703.
- [104] Kelsh RN, Eisen JS. The zebrafish colourless gene regulates development of non-ectomesenchymal neural crest derivatives. *Development* 2000;127:515–25.
- [105] Aoki Y, Saint-Germain N, Gyda M, Magner-Fink E, Lee YH, Credidio C, et al. Sox10 regulates the development of neural crest-derived melanocytes in *Xenopus*. *Dev Biol* 2003;259:19–33.
- [106] Dutton K, Dutton JR, Pauliny A, Kelsh RN. A morpholino phenotype of the colourless mutant. *Genesis* 2001;30:188–9.
- [107] Dutton KA, Pauliny A, Lopes SS, Elworthy S, Carney TJ, Rauch J, et al. Zebrafish colourless encodes sox10 and specifies non-ectomesenchymal neural crest fates. *Development* 2001;128:4113–25.
- [108] McKeown SJLV, Bronner-Fraser M, Newgreen DF, Farlie PG. Sox10 overexpression induces neural crest-like cells from all dorsoventral levels of the neural tube but inhibits differentiation. *Dev Dyn* 2005;233:430–44.
- [109] Spokony RF, Aoki Y, Saint-Germain N, Magner-Fink E, Saint-Jeannet JP. The transcription factor Sox9 is required for cranial neural crest development in *Xenopus*. *Development* 2002;129:421–32.
- [110] Cheung M, Briscoe J. Neural crest development is regulated by the transcription factor Sox9. *Development* 2003;130:5681–93.
- [111] Yan YL, Willoughby J, Liu D, Crump JG, Wilson C, Miller CT, et al. A pair of Sox: distinct and overlapping functions of zebrafish sox9 co-orthologs in craniofacial and pectoral fin development. *Development* 2005;132:1069–83.
- [112] Perez-Alcala S, Nieto MA, Barbas JA. LSox5 regulates RhoB expression in the neural tube and promotes generation of the neural crest. *Development* 2004;131:4455–65.
- [113] Southard-Smith EM, Kos L, Pavan WJ. Sox10 mutation disrupts neural crest development in *Dom* Hirshsprung mouse. *Nat Genet* 1998;18:60–4.
- [114] Kapur RP. Early death of neural crest cells is responsible for total enteric aganglionosis in Sox10<sup>Dom</sup>Sox10<sup>Dom</sup> mouse embryos. *Pediatr Dev Pathol* 1999;2:559–69.
- [115] Sonnenberg-Riethmacher E, Mische M, Stolt CC, Goerich DE, Wegner M, Riethmacher D. Development and degeneration of dorsal root ganglia in the absence of the HMG-domain transcription factor Sox10. *Mech Dev* 2001;109:253.
- [116] Gitelman I. Twist protein in mouse embryogenesis. *Dev Biol* 1997;189:205–14.
- [117] Soo K, O'Rourke MP, Khoo PL, Steiner KA, Wong N, Behringer RR, et al. Twist function is required for the morphogenesis of the cephalic neural tube and the differentiation of the cranial neural crest cells in the mouse embryo. *Dev Biol* 2002;247:251–70.
- [118] Ota M, Loebel D, O'Rourke M, N W, B T, PP T. Twist is required for patterning the cranial nerves and maintaining the viability of mesodermal cells. *Dev Dyn* 2004;230(June (2)):216–28.
- [119] Labosky PA, Kaestner KH. The winged helix transcription factor Hfh2 is expressed in neural crest and spinal cord during mouse development. *Mech Dev* 1998;76:185–90.
- [120] Odenthal J, Nusslein-Volhard C. Fork head domain genes in zebrafish. *Dev Genes Evol* 1998;208:245–58.
- [121] Kos R, Reedy MV, Johnson RL, Erickson CA. The winged-helix transcription factor FoxD3 is important for establishing the neural crest lineage and repressing melanogenesis in avian embryos. *Development* 2001;128:1467–79.
- [122] Dottori M, Gross MK, Labosky P, Goulding M. The winged-helix transcription factor Foxd3 suppresses interneuron differentiation and promotes neural crest cell fate. *Development* 2001;128:4127–38.
- [123] Meulemans D, Bronner-Fraser M. Gene-regulatory interactions in neural crest evolution and development. *Dev Cell* 2004;7:291–9.